

# You and Your Anaesthetic

**Anaesthetics**

**Surgery and Critical Care**

**Deputy Chief Executive's Office & Directorate of Operations**

**This leaflet has been designed to  
provide information to help patients  
prepare for an anaesthetic.**

# Information for patients and visitors

## Introduction

This leaflet gives basic information to help you prepare for your anaesthetic. It has been written by patients, patient representatives and anaesthetists, working in partnership.

You can find more information in other leaflets in the series on the website [www.rcoa.ac.uk/patientinfo](http://www.rcoa.ac.uk/patientinfo)

### The series also includes the following:

- Anaesthesia explained (a more detailed booklet)
- Your child's general anaesthetic
- Your spinal anaesthetic
- Epidurals for pain relief after surgery
- Headache after an epidural or spinal anaesthetic
- Your child's general anaesthetic for dental treatment
- Local anaesthesia for your eye operation
- Your tonsillectomy as day surgery
- Your anaesthetic for aortic surgery
- Anaesthetic choices for hip and knee replacement

Throughout this booklet we use the following symbols.



To highlight your options or choices.



To highlight where you may want to take particular action.



To point you to more information.

## Some Types of Anaesthesia

Anaesthesia stops you feeling pain and other sensations.

It can be given in various ways and does not always make you unconscious.

**Local anaesthesia** involves injections which numb a small part of your body. You stay conscious but free from pain.

**Regional anaesthesia** involves injections which numb a larger or deeper part of the body.

You stay conscious but free from pain

**General anaesthesia** gives a state of controlled unconsciousness.

It is essential for some operations.

You are unconscious and feel nothing.


## Anaesthetists

**Anaesthetists are doctors with specialist training who:**

- Make your experience as calm and pain free as possible
- Discuss types of anaesthesia with you and find out what you would like, helping you to make choices.
- Discuss the risks of anaesthesia with you
- Agree a plan with you for your anaesthetic and pain control
- Are responsible for giving your anaesthetic and for your well-being and safety throughout your surgery
- Manage any blood transfusions you may need
- Plan your care, if needed, in the Intensive Care Unit (ITU)

## Information for patients and visitors

### Before Coming to Hospital




Here are some things that you can do to prepare yourself for your operation.

- **If you smoke**, giving up for several weeks before the operation reduces the risk of breathing problems and makes your anaesthetic safer. The longer you can give up beforehand, the better. If you cannot stop smoking completely, cutting down will help
- **If you are very overweight**, reducing your weight will reduce many of the risks of having an anaesthetic
- **If you have loose teeth or crowns**, treatment from your dentist may reduce the risk of damage to your teeth if the anaesthetist needs to put a tube in your throat to help you breathe
- **If you have a long-standing medical problem** such as diabetes, asthma or bronchitis, thyroid problems, heart problems or high blood pressure (hypertension) you should ask your GP if you need a check-up

### Before Your Anaesthetic


You will be asked some questions to check your health before your operation. This may be at a pre-assessment clinic, by filling in a questionnaire, by talking to doctors on the ward, or when you meet your anaesthetist.



It is important for you to bring a list of:


- All the **pills, medicines, herbal remedies or supplements** you are taking, both prescribed and those that you have purchased over the counter
- Any **allergies** you may have

### On the Day of Your Operation




**Nothing to eat or drink- fasting ('Nil by Mouth')**

The hospital should give you clear instructions about fasting. It is important to follow these. If there is food or liquid in your stomach during your anaesthetic, it could come up to the back of your throat and damage your lungs.



**If you are a smoker** you should not smoke on the day of your operation. This will help reduce breathing problems during your operation.



**If you are taking medicines**, you should continue to take them as usual, unless your anaesthetist or surgeon has asked you not to. For example, if you take drugs to stop you getting blood clots (anticoagulants), aspirin drugs for diabetes or herbal remedies, you will need specific instructions.



**If you feel unwell** when you are due to come into hospital, please telephone the ward for advice.

**Your anaesthetist will meet you before your operation** and will:

- **Ask you** about your health
- **Discuss with you** which types of anaesthetic can be used
- **Discuss with you** the benefits, risks and your preferences
- **Decide with you** which anaesthetic would be best for you
- **Decide for you**, if you would prefer that

## Information for patients and visitors



**Nothing will happen to you until you understand and agree** with what has been planned for you. You have the right to refuse if you do not want the treatment suggested or if you want more information or more time to decide.



**The choice of anaesthetic depends on:**

- Your operation
- Your answers to the questions you have been asked
- Your physical condition
- Your preferences and the reasons for them
- Your anaesthetist's recommendations for you and the reasons for them
- The equipment, staff and other resources at your hospital

**Premedication** (a 'premed') is the name for drugs which are sometimes given before an anaesthetic. Some premeds prepare your body for the anaesthetic, others help you relax. They may make you more drowsy after the operation. If you want to go home on the same day, this may be delayed. If you think a premed would help you, ask your anaesthetist.

### **A needle may be used to start your anaesthetic**

If this worries you, you can ask to have a local anaesthetic cream to put on your arm to numb the skin before you leave the ward. The ward nurses should be able to do this for you.

**If you are having a local or regional anaesthetic**, you will also need to decide whether you would prefer to:

- Be fully alert
- Be relaxed and sleepy (**sedation**)
- Have a general anaesthetic as well

**Sedation** is the use of small amounts of anaesthetic or similar drugs to produce a 'sleepy-like' state.

### **When You Are Called for Your Operation**

- A member of staff will go with you to the theatre.
- A relative may be able to go with you to the theatre reception, with the exception of those admitted to the Day Surgery Unit at Grimsby. A parent will normally go with a child.
- You can wear **glasses, hearing aids and dentures** until you are in the anaesthetic room. If you are having local or regional anaesthetic, you may keep them on.
- **Jewellery and decorative piercing** should ideally be removed. If you cannot remove your jewellery, it can be covered with tape to prevent damage to it or your skin.



If you are having a local or regional anaesthetic, you can take a **personal tape, CD player or MP3 player** with you to listen to music through your headphones.



**Most people walk to theatre, others on a bed or trolley.** If you are walking, you will need your dressing gown and slippers.

# Information for patients and visitors

## The Operating Department (‘Theatres’)

Theatre staff will check your identification bracelet, your name and date of birth, and will ask you about other details in your medical records as a final check that you are having the right operation.

Your anaesthetic may start in the anaesthetic room or theatre. The anaesthetist will attach machines which measure your heart rate, blood pressure and oxygen levels.

For many anaesthetics, including some types of local anaesthetic, a needle is used to put a cannula (thin plastic tube) into a vein in the back of your hand.

## Local and Regional Anaesthetics

- Your anaesthetist will ask you to keep quite still while the injections are given
- You may notice a warm tingling feeling as the anaesthetic begins to take effect
- Your operation will only go ahead when you and your anaesthetist are sure that the area is numb
- If you are not having sedation you will remain alert and aware of your surroundings. A screen shields the operating site, so you will not see the operation unless you want to
- Your anaesthetist is always near to you and you can speak to him or her whenever you want to

## General Anaesthetics

- Anaesthetic drugs may be injected into the vein through the cannula (this is generally used for adults)

- You can breathe anaesthetic gases and oxygen through a mask, which you may hold if you prefer.


Once you are unconscious, an anaesthetist stays with you at all times and continues to give you drugs to keep you anaesthetised.

As soon as the operation is finished, the drugs will be stopped or reversed so that you regain consciousness.

After the operation, you may be taken to the **recovery room**. Recovery staff will be with you at all times. When they are satisfied that you have recovered safely from your anaesthetic you will be taken back to the ward.

## Pain Relief Afterwards

Good pain relief is important and some people need more pain relief than others. It is much easier to relieve pain if it is dealt with **before** it gets bad. Pain relief can be increased, given more often, or given in different combinations.



Occasionally, pain is a warning sign that all is not well, so you should ask for help when you feel pain.

Here are some ways of giving pain relief:

- **Pills, tablets or liquids to swallow**

These are used for all types of pain. They take at least half an hour to work. You need to be able to eat, drink and not feel sick for these drugs to work.

- **Injections**

These are often needed, and are given either into a vein (through your cannula) for a quicker effect, or into your leg or buttock muscle. If in a muscle they may take up to 20 minutes to work.



# Information for patients and visitors

- **Suppositories**

These waxy pellets are put in your back passage (rectum). The pellet dissolves and the drug passes into the body. They are useful if you cannot swallow or if you might vomit.

- **Patient-controlled analgesia (PCA)**

This is a method using a machine that allows you to control your pain relief yourself. If you would like more information ask for a leaflet on PCA.



## Local anaesthetics and regional blocks

These types of anaesthesia can be very useful for relieving pain after surgery. More details can be found in the leaflet 'Epidurals for pain relief after surgery' or ask your Anaesthetists for options available locally.

## What will I feel like afterwards?

How you feel will depend on the type of anaesthetic and operation you have had, how much pain relieving medicine you need and your general health.

## Understanding Risk

In modern anaesthesia, serious problems are uncommon. Risk cannot be removed completely, but modern equipment, training and drugs have made it a much safer procedure in recent years.

### To understand a risk, you must know:

- How likely it is to happen
- How serious it could be
- How it can be treated

### The risk to you as an individual will depend on:

- Whether you have any other illness
- Personal factors, such as smoking or being overweight
- Surgery which is complicated, long or done in an emergency.

### People vary in how they interpret words and numbers.

This scale is provided to help:

Very Common	Common	Uncommon	Rare	Very Rare
1 in 10	1 in 100	1 in 1000	1 in 10,000	1 in 100,000

## Side Effects and Complications



More information about risks associated with having an anaesthetic can be found on [www.rcoa.ac.uk/patientinfo](http://www.rcoa.ac.uk/patientinfo) More information on the side effects and complications listed here is in the booklet 'Anaesthesia explained'.

RA = May occur with a regional anaesthetic

GA = May occur with a general anaesthetic.

### Very common and common side effects

- |    |    |                                          |
|----|----|------------------------------------------|
| RA | GA | Feeling sick and vomiting after surgery  |
|    | GA | Sore throat                              |
| RA | GA | Dizziness, blurred vision                |
| RA | GA | Headache                                 |
| RA | GA | Bladder problems                         |
|    | GA | Damage to lips or tongue (usually minor) |
| RA | GA | Itching                                  |
| RA | GA | Aches, pains and backache                |



## Information for patients and visitors

- RA GA Pain during injection of drugs
- RA GA Bruising and soreness
- GA Confusion or temporary memory loss

### Uncommon side effects and complications

- GA Chest infection
- GA Muscle pains
- RA GA Slow breathing (depressed respiration)
- GA Damage to teeth
- RA GA An existing medical condition getting worse
- GA Awareness (becoming conscious during your operation)

### Rare or very rare complications

- GA Damage to the eyes
- RA GA Heart attack or stroke
- RA GA Serious allergy to drugs
- RA GA Nerve damage
- RA GA Death
- RA GA Equipment failure

Deaths caused by anaesthesia are very rare, and are usually caused by a combination of 4 or 5 complications together. There are probably about five deaths for every million anaesthetics in the UK.

### Questions You May Like to Ask Your Anaesthetist

- Q Who will give my anaesthetic?
- Q Do I have to have a general anaesthetic?
- Q What type of anaesthetic do you recommend?

- Q Have you often used this type of anaesthetic?
- Q What are the risks of this type of anaesthetic?
- Q Do I have any special risks?
- Q How will I feel afterwards?

### Tell Us What You Think



The Association of Anaesthetists of Great Britain and Ireland (AAGBI)



The Royal College of Anaesthetists (RCoA)

© The RCoA and AAGBI agree to the copying of this document for the purpose of producing local leaflets in the United Kingdom and Ireland. This information was adapted from a patient information leaflet template on the AAGBI and RCoA websites. The Patient Information Unit must agree to any changes if the AAGBI and RCoA crests are to be kept.

This publication includes text taken from the Royal College of Anaesthetists' (RCoA) leaflet but the RCoA has not reviewed this as a whole.

# Information for patients and visitors

---

## Concerns and Queries

If you have any concerns / queries about any of the services offered by the Trust, in the first instance, please speak to the person providing your care.

### For Diana, Princess of Wales Hospital

Alternatively you can contact the Patient Advice and Liaison Service (PALS) on (01472) 875403 or at the PALS office which is situated near the main entrance.

### For Scunthorpe General Hospital

Alternatively you can contact the Patient Advice and Liaison Service (PALS) on (01724) 290132 or at the PALS office which is situated on C Floor.

Alternatively you can email:  
[nlq-tr.PALS@nhs.net](mailto:nlq-tr.PALS@nhs.net)

## Northern Lincolnshire and Goole NHS Foundation Trust

**Diana Princess of Wales Hospital**  
Scartho Road  
Grimsby  
DN33 2BA  
01472 874111

**Scunthorpe General Hospital**  
Cliff Gardens  
Scunthorpe  
DN15 7BH  
01724 282282

**Goole & District Hospital**  
Woodland Avenue  
Goole  
DN14 6RX  
01405 720720

[www.nlg.nhs.uk](http://www.nlg.nhs.uk)

Date of issue: December, 2016

Review Period: December, 2019

Author: Consultant Anaesthetics

IFP-265 v4.2

© NLGHT 2016

